

http://didawiki.di.unipi.it/doku.php/ magistraleinformatica/psc/start

PSC 2021/22 (375AA, 9CFU)

Principles for Software Composition

Roberto Bruni http://www.di.unipi.it/~bruni/

15 - HOFL: Consistency?

HOFL Operational vs Denotational

Differences

operational $t \rightarrow c$

closed, typeable terms no environment not a congruence canonical terms denotational $[t] \rho$

typeable terms environment congruence mathematical entities

$$\forall t, c. \quad t \to c \quad \stackrel{?}{\Leftrightarrow} \quad \forall \rho. \ [t] \rho = [c] \rho$$
$$t \to c \Rightarrow \forall \rho. \ [t] \rho = [c] \rho$$
$$(\forall \rho. \ [t] \rho = [c] \rho) \neq t \to c \quad \text{there is on}$$

there is only one type for which the implication holds

Inconsistency: example

x: int $c_0 = \lambda x. x + 0$ $c_1 = \lambda x. x$

already in canonical forms


 $\llbracket c_0 \rrbracket \rho = \llbracket c_1 \rrbracket \rho \qquad \qquad c_0 \not\rightarrow c_1$

 $\llbracket c_0 \rrbracket \rho = \llbracket \lambda x. x + 0 \rrbracket \rho = \lfloor \lambda d. d + \lfloor 0 \rfloor \rfloor = \lfloor \lambda d. d \rfloor = \llbracket \lambda x. x \rrbracket \rho = \llbracket c_1 \rrbracket \rho$

Correctness $t \to c \Rightarrow \forall \rho. \ [t] \rho = [c] \rho$

proof. we proceed by rule induction

$$P(t \to c) \stackrel{\text{def}}{=} \forall \rho. \ [t] \rho = [c] \rho$$



$$P(c \rightarrow c) \stackrel{\text{def}}{=} \forall \rho. \ [c] \rho = [c] \rho$$
 obvious

$t \to c \Rightarrow \forall \rho. \ [t] \rho = [c] \rho$

(continue)

$$\frac{t_1 \to n_1 \quad t_2 \to n_2}{t_1 \text{ op } t_2 \to n_1 \text{ op } n_2}$$
 assume
$$P(t_1 \to n_1) \stackrel{\text{def}}{=} \forall \rho. \ [t_1] \rho = [n_1] \rho = \lfloor n_1 \rfloor$$
$$P(t_2 \to n_2) \stackrel{\text{def}}{=} \forall \rho. \ [t_2] \rho = [n_2] \rho = \lfloor n_2 \rfloor$$

we prove $P(t_1 \text{ op } t_2 \to n_1 \text{ op } n_2) \stackrel{\text{def}}{=} \forall \rho. [t_1 \text{ op } t_2] \rho = [n_1 \text{ op } n_2] \rho$

$$\begin{bmatrix} t_1 \text{ op } t_2 \end{bmatrix} \rho = \begin{bmatrix} t_1 \end{bmatrix} \rho \underbrace{\text{op}}_{\perp} \begin{bmatrix} t_2 \end{bmatrix} \rho$$
$$= \begin{bmatrix} n_1 \end{bmatrix} \underbrace{\text{op}}_{\perp} \begin{bmatrix} n_2 \end{bmatrix}$$
$$= \begin{bmatrix} n_1 \underbrace{\text{op}} n_2 \end{bmatrix}$$
$$= \begin{bmatrix} n_1 \underbrace{\text{op}} n_2 \end{bmatrix} \rho$$

(by definition of $\llbracket \cdot \rrbracket$) (by inductive hypotheses) (by definition of \underline{op}_{\perp}) (by definition of $\llbracket \cdot \rrbracket$)

$t \to c \Rightarrow \forall \rho. \ \llbracket t \rrbracket \rho = \llbracket c \rrbracket \rho$

(continue)

$$t \to 0 \quad t_0 \to c_0$$

if t then t_0 else $t_1 \rightarrow c_0$

assume

$$P(t \to 0) \stackrel{\text{def}}{=} \forall \rho. \ [t] \rho = [0] \rho = [0]$$

$$P(t_0 \to c_0) \stackrel{\text{def}}{=} \forall \rho. \ [t_0] \rho = [c_0] \rho$$

We prove $P(\text{if } t \text{ then } t_0 \text{ else } t_1 \rightarrow c_0) \stackrel{\text{def}}{=} \forall \rho$. [[if $t \text{ then } t_0 \text{ else } t_1]] <math>\rho = [[c_0]] \rho$

$$\begin{bmatrix} \text{if } t \text{ then } t_0 \text{ else } t_1 \end{bmatrix} \rho = Cond(\llbracket t \rrbracket \rho, \llbracket t_0 \rrbracket \rho, \llbracket t_1 \rrbracket \rho) & \text{(by def. of } \llbracket \cdot \rrbracket) \\ = Cond(\lfloor 0 \rfloor, \llbracket t_0 \rrbracket \rho, \llbracket t_1 \rrbracket \rho) & \text{(by ind. hyp.)} \\ = \llbracket t_0 \rrbracket \rho & \text{(by def. of } Cond) \\ = \llbracket c_0 \rrbracket \rho & \text{(by ind. hyp.)} \end{aligned}$$

ifn) analogous (omitted)

$t \to c \Rightarrow \forall \rho. \ \llbracket t \rrbracket \rho = \llbracket c \rrbracket \rho$

(continue)

$$\frac{t \to (t_0, t_1) \quad t_0 \to c_0}{\mathbf{fst}(t) \to c_0}$$

assume

$$P(t \to (t_0, t_1)) \stackrel{\text{def}}{=} \forall \rho. \ [t] \rho = [(t_0, t_1)] \rho$$

$$P(t_0 \to c_0) \stackrel{\text{def}}{=} \forall \rho. \ [t_0] \rho = [c_0] \rho$$

we prove
$$P(\mathbf{fst}(t) \to c_0) \stackrel{\text{def}}{=} \forall \rho. [\![\mathbf{fst}(t)]\!] \rho = [\![c_0]\!] \rho$$

$$\llbracket \mathbf{fst}(t) \rrbracket \rho = \pi_1^*(\llbracket t \rrbracket \rho) \qquad \text{(by def. of } \llbracket \cdot \rrbracket) \\ = \pi_1^*(\llbracket (t_0, t_1) \rrbracket \rho) \qquad \text{(by ind. hyp.)} \\ = \pi_1^*(\lfloor (\llbracket t_0 \rrbracket \rho, \llbracket t_1 \rrbracket \rho) \rfloor) \qquad \text{(by def. of } \llbracket \cdot \rrbracket) \\ = \pi_1(\llbracket t_0 \rrbracket \rho, \llbracket t_1 \rrbracket \rho) \qquad \text{(by def. of lifting)} \\ = \llbracket t_0 \rrbracket \rho \qquad \text{(by def. of } \pi_1) \\ = \llbracket c_0 \rrbracket \rho \qquad \text{(by ind. hyp.)}$$

snd) analogous (omitted)

$t \to c \Rightarrow \forall \rho. \ \llbracket t \rrbracket \rho = \llbracket c \rrbracket \rho$

(continue)

$$\frac{t_1 \to \lambda x. t_1' \quad t_1' [t_0/x] \to c}{(t_1 \ t_0) \to c}$$

assume

$$P(t_1 \to \lambda x. t_1') \stackrel{\text{def}}{=} \forall \rho. [[t_1]] \rho = [[\lambda x. t_1']] \rho$$

$$P(t_1'[t_0/x] \to c) \stackrel{\text{def}}{=} \forall \rho. [[t_1'[t_0/x]]] \rho = [[c]] \rho$$

we prove $P((t_1 t_0) \rightarrow c) \stackrel{\text{def}}{=} \forall \rho . [[(t_1 t_0)]] \rho = [[c]] \rho$ $\|(t_1 t_0)\| \rho = \mathbf{let} \ \varphi \leftarrow \|t_1\| \rho. \ \varphi(\|t_0\| \rho)$ (by definition of $\|\cdot\|$) = let $\varphi \leftarrow [\lambda x. t'_1] \rho. \varphi([t_0] \rho)$ (by ind. hypothesis) = let $\varphi \leftarrow |\lambda d. [t_1] \rho[d/x] | . \varphi([t_0] \rho)$ (by definition of $[\cdot]$) = $(\lambda d. [t_1'] \rho [d/x]) ([t_0] \rho)$ (by de-lifting) $= \llbracket t_1' \rrbracket \rho [\llbracket t_0 \rrbracket \rho / _x]$ (by application) $= [t'_1[t_0/x]] \rho$ (by Subst. Lemma) $= \llbracket c \rrbracket \rho$ (by ind. hypothesis)

t^{rec}

$$t \to c \Rightarrow \forall \rho. \ [t] \ \rho = [c] \ \rho$$
 (continue)

$$\left| \frac{x \cdot t}{x} \right| \to c$$
 ass

rec *x*. $t \rightarrow c$

assume

 $P(t[^{\mathbf{rec}\ x.\ t}/_{x}] \to c) \stackrel{\text{def}}{=} \forall \rho. \ \left[\!\left[t[^{\mathbf{rec}\ x.\ t}/_{x}]\right]\!\right] \rho = \left[\!\left[c\right]\!\right] \rho$

we prove $P(\operatorname{rec} x. t \to c) \stackrel{\text{def}}{=} \forall \rho. [[\operatorname{rec} x. t]] \rho = [[c]] \rho$

 $\llbracket \operatorname{rec} x. t \rrbracket \rho = \llbracket t \rrbracket \rho \llbracket \operatorname{rec} x. t \rrbracket \rho /_{x} \end{bmatrix}$ $= \llbracket t \llbracket \operatorname{rec} x. t /_{x} \rrbracket \rho$ $= \llbracket c \rrbracket \rho$

(by definition)(by the Substitution Lemma)(by inductive hypothesis)

HOFL convergence Operational vs Denotational

Operational convergence

 $t: \tau$ closed

 $t \downarrow \quad \Leftrightarrow \quad \exists c \in C_{\tau}. \ t \longrightarrow c$ $t \uparrow \quad \Leftrightarrow \quad \neg \ t \downarrow$

Examples $\mathbf{rec} x. x \uparrow$ $\lambda y. \mathbf{rec} x. x \downarrow$ $(\lambda y. \mathbf{rec} x. x) 0 \uparrow$ if 0 then 1 else rec $x. x \downarrow$

Denotational converg.

 $t:\tau$ closed

 $t \Downarrow \Leftrightarrow \forall \rho \in Env, \exists v \in V_{\tau}. \llbracket t \rrbracket \rho = \lfloor v \rfloor$

 $t \Uparrow \quad \Leftrightarrow \quad \neg \quad t \Downarrow$

Examples $\begin{bmatrix} \operatorname{rec} x. x \end{bmatrix} \rho \Uparrow$ $\begin{bmatrix} \lambda y. \operatorname{rec} x. x \end{bmatrix} \rho \Downarrow$ $\begin{bmatrix} (\lambda y. \operatorname{rec} x. x) \ 0 \end{bmatrix} \rho \Uparrow$ $\begin{bmatrix} \operatorname{if} 0 \text{ then } 1 \text{ else rec } x. x \end{bmatrix} \rho \Downarrow$

Consistency on converg.		
TH. $t: \tau$ closed $t \downarrow \Rightarrow t \Downarrow$		
proof. $t\downarrow$	$\Rightarrow t \rightarrow c$	by def (for some c)
	$\Rightarrow \forall \rho. \llbracket t \rrbracket \rho = \llbracket c \rrbracket \rho$	by correctness
	$\Rightarrow \forall \rho. \llbracket t \rrbracket \rho \neq \bot$	canonical $\llbracket c \rrbracket \rho \neq \bot$
	$\Rightarrow t \Downarrow$	by def
TH. $t:\tau$ close	sed $t \Downarrow \Rightarrow t \downarrow$	

the proof is not part of the program of the course (structural induction would not work)

HOFL equivalence Operational vs Denotational

HOFL equivalences

 $t_0, t_1: \tau$ closed

 $t_0 \equiv_{\text{op}} t_1 \quad \text{iff} \quad \forall c. \ t_0 \to c \iff t_1 \to c$

 $t_0 \equiv_{\text{den}} t_1$ iff $\forall \rho$. $\llbracket t_0 \rrbracket \rho = \llbracket t_1 \rrbracket \rho$

Op is more concrete

TH. $\equiv_{op} \subseteq \equiv_{den}$

proof. take $t_0, t_1 : \tau$ closed, such that $t_0 \equiv_{op} t_1$

either $\exists c. t_0 \rightarrow c \land t_1 \rightarrow c \text{ or } t_0 \uparrow \land t_1 \uparrow$

if $\exists c. t_0 \rightarrow c \land t_1 \rightarrow c$

by correctness $\forall \rho$. $\llbracket t_0 \rrbracket \rho = \llbracket c \rrbracket \rho = \llbracket t_1 \rrbracket \rho$ thus $t_0 \equiv_{den} t_1$

if
$$t_0 \uparrow \land t_1 \uparrow$$

by agreement on convergence $t_0 \Uparrow \land t_1 \Uparrow$

i.e.
$$\forall \rho$$
. $\llbracket t_0 \rrbracket \rho = \bot_{D_\tau} = \llbracket t_1 \rrbracket \rho$ thus $t_0 \equiv_{\text{den}} t_1$

Den is strictly more abstract

TH.
$$\equiv_{den} \not\subseteq \equiv_{op}$$

proof.

see previous counterexample

x: *int* $c_0 = \lambda x. x + 0$ $c_1 = \lambda x. x$

Consistency on int

TH. t: int closed $t \to n \quad \Leftrightarrow \quad \forall \rho. \llbracket t \rrbracket \rho = \lfloor n \rfloor$

proof.

 $\Rightarrow) \text{ if } t \rightarrow n \text{ then } \llbracket t \rrbracket \rho = \llbracket n \rrbracket \rho = \lfloor n \rfloor$

 $\Leftarrow) \text{ if } \llbracket t \rrbracket \rho = \lfloor n \rfloor \text{ it means } t \Downarrow \\ \text{ by agreement on convergence } t \downarrow \\ \text{ thus } t \to m \text{ for some } m \\ \text{ but then by correctness } \llbracket t \rrbracket \rho = \llbracket m \rrbracket \rho = \lfloor m \rfloor \\ \text{ and it must be } m = n \end{cases}$

Equivalence on int

TH.
$$t_0, t_1 : int$$
 $t_0 \equiv_{op} t_1 \Leftrightarrow t_0 \equiv_{den} t_1$

proof. we know $t_0 \equiv_{op} t_1 \Rightarrow t_0 \equiv_{den} t_1$ we prove $t_0 \equiv_{den} t_1 \Rightarrow t_0 \equiv_{op} t_1$

assume $t_0 \equiv_{den} t_1$ either $\forall \rho$. $\llbracket t_0 \rrbracket \rho = \bot_{\mathbb{Z}_\perp} = \llbracket t_1 \rrbracket \rho$ or $\forall \rho$. $\llbracket t_0 \rrbracket \rho = \lfloor n \rfloor = \llbracket t_1 \rrbracket \rho$ for some n

if $\forall \rho$. $\llbracket t_0 \rrbracket \rho = \bot_{\mathbb{Z}_{\perp}} = \llbracket t_1 \rrbracket \rho$ then $t_0 \Uparrow, t_1 \Uparrow$

by agreement on convergence $t_0 \uparrow, t_1 \uparrow$ thus $t_0 \equiv_{op} t_1$

if
$$\forall \rho$$
. $\llbracket t_0 \rrbracket \rho = \lfloor n \rfloor = \llbracket t_1 \rrbracket \rho$ then $t_0 \to n$, $t_1 \to n$
thus $t_0 \equiv_{\text{op}} t_1$

HOFL Unlifted Semantics

Unlifted Domains

 $D_{\tau} \triangleq (V_{\tau})_{\perp} \qquad \text{lifted domains}$ $V_{int} \triangleq \mathbb{Z}$ $V_{\tau_1 * \tau_2} \triangleq D_{\tau_1} \times D_{\tau_2} = (V_{\tau_1})_{\perp} \times (V_{\tau_2})_{\perp}$ $V_{\tau_1 \to \tau_2} \triangleq [D_{\tau_1} \to D_{\tau_2}] = [(V_{\tau_1})_{\perp} \to (V_{\tau_2})_{\perp}]$

unlifted domains

$$U_{int} \triangleq \mathbb{Z}_{\perp}$$
$$U_{\tau_1 * \tau_2} \triangleq U_{\tau_1} \times U_{\tau_2}$$
$$U_{\tau_1 \to \tau_2} \triangleq [U_{\tau_1} \to U_{\tau_2}]$$

Unlifted Semantics as before $(n) \rho \triangleq |n|$ $(x) \rho \triangleq \rho(x)$ $(t_1 \text{ op } t_2) \rho \triangleq (t_1) \rho \text{ op} (t_2) \rho$ (if t then t_1 else t_2) $\rho \triangleq \operatorname{Cond}_{\tau}(\langle t \rangle \rho, \langle t_1 \rangle \rho, \langle t_2 \rangle \rho)$ $(|\mathbf{rec} x. t|) \rho \triangleq fix \lambda d. (|t|) \rho |d|_x$ $((t_1, t_2))\rho \triangleq ((t_1)\rho, (t_2)\rho)$ without lifting $(\mathbf{fst}(t)) \rho \triangleq \pi_1 ((t) \rho)$ $(|\mathbf{snd}(t)|) \rho \triangleq \pi_2 (||t|) \rho$ $(\lambda x. t) \rho \triangleq \lambda d. (t) \rho [d/x]$ $(t t_0) \rho \triangleq ((t) \rho) ((t_0) \rho)$

$$\begin{aligned} & \underbrace{t_{1} \triangleq \operatorname{rec} x. \ x \ : int \to int}_{x: \ int \to int} \quad \underbrace{t_{2} \triangleq \lambda y. \ \operatorname{rec} z. \ z \ : int \to int}_{y, z: \ int} \\ & \underbrace{D_{int \to int} = [\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}]_{\perp}}_{\left[t_{1}]\right] \rho = \perp_{\left[\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}\right] \perp} \quad \left[t_{2}\right] \rho = \lfloor \perp_{\left[\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}\right]} \right] \\ & \underbrace{t_{1} \uparrow}_{t_{1}} \quad \underbrace{t_{2} \downarrow}_{t_{2} \to t_{2}} \\ & \underbrace{U_{int \to int} = [\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}]}_{\left[t_{1}\right] \rho = \perp_{\left[\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}\right]} \quad \left[t_{2}\right] \rho = \perp_{\left[\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}\right]} \\ & \underbrace{t_{1} \uparrow}_{t_{1}} \quad \underbrace{t_{2} \downarrow}_{t_{2} \to t_{2}} \\ & \underbrace{U_{int \to int} = [\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}]}_{t_{1}} \quad \underbrace{t_{2} \downarrow \rho = \perp_{\left[\mathbb{Z}_{\perp} \to \mathbb{Z}_{\perp}\right]} = \lambda d. \perp_{\mathbb{Z}_{\perp}} \\ & \underbrace{t_{2} \downarrow \neq t_{2} \downarrow_{unlifted}}_{t_{2} \downarrow \neq t_{2} \downarrow_{unlifted}} \end{aligned}$$